



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

bacter, particularly in the présence of cellulose decomposing organisms.

The high acidity of the cranberry soils would preclude the very idea of finding the *Azotobacter* in these soils and the early students⁶ of this group of organisms were of the opinion that they can not live in acid media at all but the reaction has to be adjusted first to neutrality before the conditions are made favorable for their activities.

A Savannah bottom cranberry bog situated at Whitesbog, N. J., was used for this work. A part of the bog was limed three years ago and the crop was almost double of the corresponding plot, unlimed. Samples of the soil from the two plots were secured under sterile conditions and used for this study. The soil is nothing more than some white sand interwoven with decayed and living plant residues.

The hydrogen-ion concentration of the two soils was determined by means of the colorimetric method, using the phenol-sulfon-phthalein indicators suggested by Clark and Lubs.⁷ The method corresponds very closely with the electrometric determinations using the hydrogen electrode, as was shown by Gillespie.² A definite amount of soil was shaken with double its weight of distilled water, then centrifuged; the supernatant clear liquid was syphoned off and used for the determination of the hydrogen-ion concentration. The unlimed soil had an hydrogen-ion concentration of $\text{pH} = 5.4$ to 5.6 while for the limed soil pH was $6.2-6.4$.

The two soils were added in 10-gram quantities to 100 cc.c. portions of a sterile faintly alkaline nitrogen-free mannite solution and incubated at 25° . The solution in the flasks containing the limed soils became turbid in four days and a pellicle characteristic of *Azotobacter* began to develop in some flasks. On microscopic examination the solution was found to contain an abundance of *Azotobacter* cells and *Actinomyces* filaments. The solution in all the flasks to which the unlimed soil was added remained clear as in the control, but has shown a profuse gas production.

⁶ Lipman, *Ann. Rept. N. J. Agr. Exp. Sta.*, pp. 262-268, 1904.

⁷ *Jour. Bact.*, Vol. 2, Nos. 1, 2, 3, 1917.

On microscopic examination no *Azotobacter* cells and no *Actinomyces* filaments were discovered.

The limiting reaction for the existence of *Azotobacter* in the soil, expressed in the hydrogen-ion concentration is thus found to fall between $\text{pH} = 5.4$ to 5.6 and $\text{pH} = 6.2$ to 6.4 and is probably nearer the latter. This will confirm the results of Gainey¹ and Christensen³ that an hydrogen-ion concentration of the soil $= \text{pH} = 6.0$ is the limiting reaction for the activities of *Azotobacter* in the soil.

The occurrence of *Actinomyces* filaments together with *Azotobacter* cells suggests a still more interesting and important possibility, association between these two groups of soil microorganisms. As will be soon shown elsewhere many *Actinomyces* decompose organic residues very rapidly. The association between these two groups of organisms, change of reaction, and the action of *Actinomyces* upon the nitrogen-fixation by *Azotobacter* is being studied at present in this laboratory.

The importance of *Azotobacter* in cranberry soils, which can be effected by changing the reaction of those soils, thus becomes apparent: these organisms, whether alone or in association with others, utilize the plant residues as a source of energy and this allows them to fix the atmospheric nitrogen and increase its supply in the soil, which goes towards an increased crop production.

SELMAN A. WAKSMAN

N. J. AGR. EXP. STATION,
NEW BRUNSWICK, N. J.,
November 2, 1918

SCIENCE

A Weekly Journal devoted to the Advancement of
Science, publishing the official notices and pro-
ceedings of the American Association for
the Advancement of Science

Published every Friday by

THE SCIENCE PRESS

LANCASTER, PA.

GARRISON, N. Y.

NEW YORK, N. Y.

Entered in the post-office at Lancaster, Pa., as second class matter